

VOL 1, PP 85-92.

May 15, 1905

INVERTEBRATA PACIFICA

EDITED BY C. F. BAKER, Estación Agronómica, Santiago de las Vegas, Cuba.

NEUROPTEROID INSECTS

NOTES ON NEUROPTEROID INSECTS OF

THE PACIFIC

COAST OF NORTH AMERICA WITH DESCRIPTIONS
OF NEW SPECIES BY NATHAN BANKS.

C. F. BAKER.

Through the kindness of Dr. Calvert and Mr. Banks I am able now to present the following preliminary report on the collections of Neuropterold Insects brought together by me during my stay on the Pacific coast.

EPHEMERIDA

3122 Callibaetis undata Pict.

Occasional at Claremont, California. Det. Banks.

2241 Cleon sp.

Taken in the mountains near Claremont, California, Det. Banks.

2242 Heptagenia n. sp.

A common form in the mountains near Claremont, California, in the Springtime, dancing in the sunbeams where they break through the thick vegetation of the deep rich gulches.

ODONATA

1549 Hetaerina americana Fab. race californica Hag. Occasional at Claremont, California. Det. Calvert. 1255 and 565 Ischnura perparva Selvs.

Common about Lagunita at Stanford University, California. Det. Calvert.

1550 Ischnura ramburi Selys var. credula Hag.

Occasional at Chinandega, Nicaragua. Det. Calvert.

1579 Leptobasis vacillans Selys.

Taken at Granada, Nicaragua, and also at Chinandega, Det. Calvert and Banks.

1256 Ceratura capreola Hag.

Several specimens were taken at Managua, Nicaragua, Det. Calvert. 586 and 566 Enallagma carunculatum Morse.

Common about Lagunita at Stanford University, California. Det. Calver.

3419 Enallagma calverti Morse.

Occasional in Ormsby county, Nevada. Det. Calvert.

580 and 1260 Argia puella Hag.

Frequent at Chinandega, Nicaragua. Det. Calvert and Banks.

1547 Argia vivida Hag.

Common at Claremont, California. Det. Calvert.

587 Lestes stulta Hag.

Abundant about Lagunita at Stanford University, California, Det. Calvert and Banks.

3624 Amphiagrion saucium Burm, race abbreviatum Selvs,

A large series taken at Pine Łake in Southern California by Mr. James Johnson, Det. Calvert.

1261 Neoneura amelia Calvert.

Occasional at Chinandega, Nicaragua. Det. Calvert.

1251 Sympetrum pallipes Hag.

Occasional at Stanford University, California, Det. Calvert.

1252 Sympetrum illotum Hag.

Occasional at Stanford University, California, Det. Calvert.

1546 Libellula saturata Uhler,

Common about Lagunita at Stanford University, California, and at Claremont. Det. Calvert.

584 Mesothemis corrupta Hag.

Frequent at Stanford University, California, Det. Calvert and Banks.

1253 Trithemis funerea Hag.

Occasional at Stanford University, California. Det. Calvert.

1545 Paltothemis lineatipes Karsch.

Occasional at Claremont, California. "Not hitherto been found in California, although it is known from Arizoua and Texas. It was originally described from Brazil and subsequently recorded from Mexico and Venezuela". (Calvert).

1544 Tramea lacerata Hag.

Occasional at Claremont, California. Det. Calvert.

1543 Aeschna multicolor Hag.

Frequent at Claremont, California, and in Ormsby county, Nevada. Det. Calvert.

PLECOPTERA

2221 Perla californica Banks n. sp.

"Head brown, pale reddish yellow between ocelli and on the clypeus; antennae pale brown; prothorax brown, with a trace of paler median line, rest of thorax dark brown, with a pale median stripe; abdomen pale brown; tibiae paler, with a blackish band on base; setae yellowish brown. Wings nearly hyaline, venation pale brownish; ventral plate of female pale, with a prominent, rounded blackish spot each side. Pronotnm plainly broader than long, anterior angles acute, posterior ones slightly rounded. Structure similar to P. lycorias. Length 30 mm. Claremont, California.

Readily separated from its allies by the two black spots on ventral plate."

3395 Nemoura depressa Banks.

Taken at light in Claremont, California. Det. Banks.

3397 and 3398 Tinodes consueta McLach.

Occasional at Claremont, California, Det. Banks,

3115 Sialis infumata Newm.

Two specimens taken at Claremont, California. Det. Banks.

ISOPTERA

112 Termopsis angusticollis Wlk.

Workers and winged males common in old logs at Claremont, California, and in Ormsby county, Nevada. Det. Banks.

CORRODENTIA

2 Elipsocus n. sp.?

Taken in great numbers from evergreen oaks very early in Spring at Stanford University, California. Det. Banks.

3 Caecilius sp. perhaps aurantiacus.

Occasional at Stanford University, California. Det. Banks.

3128 Caecilius sp.

Frequent on oaks at Claremont, California. Det. Banks.

3125 Psocus sp.

Occasional at Claremont, California, Det. Banks,

3130 Psocus californicus Banks.

Occasional at Claremont, California, but more common in the nearby mountains. Det. Banks.

3127 Peripsocus californicus Banks.

Occasional at Stanford University, California. Det. Banks.

NEUROPTERA S. STR.

All of the species cited under this head have been determined by Mr. Banks.

573 Raphidia oblita Hag.

Common at Stanford University, California.

8 Raphidia bicolor Alb.

Common in Ormsby county, Nevada.

2222 Raphidia adnixa Hag.

Common in Ormsby county, Nevada. Found also at Stanford University, California.

2126 Raphidia occulta Banks.

Common at Claremont California and in the nearby mountains,

3426 Mantispa brevicollis Banks.

Taken in Ormsby county, Nevada,

574 Mantispa viridis Wlk.

Occasional at Granada, Nicaragua.

7 Chrysopa fraterna Banks.

Common in Ormsby county, Nevada.

4 Chrysopa californica Coq.

Comon in Ormsby county, Nevada and at Stanford University,

3117 Chrysopa externa Hag.

Common at Claremont, California, and in the nearby mountains.

3120 Chrysopa bimaculata McClendon.

Occasional at Chinandega, Nicaragua.

3402 Chrysopa rufilabris Burm.

Taken at Elsinore, San Bernardino county, California,

3392 to 3394 Chrysopa coloradensis Banks.

Collected in Ormsby county, Nevada and also in the mountains near Claremont, California.

3399 Eremochrysa fraterna Banks.

Taken at Pine Lake, Southern California, by James D. Jonhson, a former student of mine.

11 Brachynemurus peregrinus Hag.

Common in Ormsby county, Nevada.

10 Brachynemurus ferox Walk.

Frequent in Ormsby county, Nevada.

2232 Brachynemurus longipalpis Hag.

Oceasional at Claremont, California.

2230 Brachynemurus Sackeni Hag.

Occasional at Claremont, California.

2233 Brachynemurus papago Currie.

Occasional at Claremont, California.

576 Ascalaphus tuberculatus Banks.

Frequent near the Indian pueblo of San Marcos in Western Nicarragua.

3405 Polystoechotes punctatus Fabr.

Mr. Culbertson, a former student of mine, took several specimens of this species at Three Rivers, California.

4035 Berotha occidentalis Banks n. sp.

"Face yellowish; vertex brown: antennae pale yellowish: thorax brown, rather densely clothed with brown bristles; abdomen pale brown, more yellowish on the Lase, legs very pale, clothed with gray hair and black bristles; wings tinted with brown, venation brown, there are many scattered round brown spots, each centered upon a vein. Along the costal area are several large dark brown spots, and the pterostigma quite dark brown; the apical margin is brown interrupted with white, gradate veinlets and the basal cross veinlets heavily dark brown. Hind wings hyaline, pterostigma brownish, and the cross vein below it nargined with brown. The wings are of the usual shape, but the outer margin is evenly and deeply concave, not all sinuate; there are five branches of the radial sector, and the latter is connected to the radius by a cross vein near the middle of the wings. The male has at the tip of the abdomen a bristly style on each side. Length to tip of wings 12 mm.

Ormsby county, Nevada, July, Baker, and also at Phoenix, Arizona (another collector).''

3422 Hemerobius coloradensis Banks.

Occasional in Ormsby county, Nevada.

3423 Hemerobius pacificus Banks.

Occasional in Ormsby county, Nevada.

3424 Hemerobius californicus Banks.

Taken at Stanford University, California.

2336 Hemerobius n. sp.

Taken at Granada, Nicaragua.

2223 Megalomus minor Banks n. sp.

"Head pale yellowish, vertex with a dark median stripe. Antennae pale, rather darker beyond the middle, palpi pale yellowish. Thorax and abdomen dark brown; legs pale yellowish. Wings hyaline, venation pale, marked with brown, the longitudinal veins and costal coinlets interruptedly brown and yellowish; the gradate series nearly black and a black spot at anal angle, and a smaller one on cubitus toward the base; margin with alternate brown and white patches. Wings of usual shape; venation not as dense as in M. moestus and M. latus: five or six sectors; the first gradate series is very regular; the outer series but little curved. Length 6 mm.

Two specimens from San Marcos, Nicaragua, and one from Ormsby county, Nevada (Baker). Differs from M. moestus and M. latus in the smaller size, the dark stripe on vertex, and the less dense venation. M. pictus Hagen from Mexico is unknown to me, but is said to have the basal joints of the antennae black, which is not the case in the other three species.'

3124 Sympherobius angustus Banks.

Occasional at Claremont, California,

3123 Micromus variolosus Hag.

Taken at Claremont, Los Angeles county, California.

3421 Micromus montanus Hag.

Occasional in Ormsby county, Nevada.

3403 Maracandula bellula Banks.

Collected for me at Three Rivers, California, by Mr. Calbertson.

575 Myrmeleon rusticus Hag.

Frequent at Granada, Nicaragua.

2228 Myrmeleon distans Banks.

Occasional at Claremont, California,

12 Myrmeleon immaculatus DeG, var coloradensis Banks.

Occasional in Ormsby county, Nevada.

MECOPTERA

583 Bittacus apterus McLachlan.
Frequent about Lagunita at Stanford University, California.

TRICHOPTERA

2235 **Leptocella** sp. Occasional at Chinandega, Nicaragua.

577 **Chimarrha** sp.

Taken at Chinandega, Nicaragna.

2237 Rhyacophila coloradensis Bank.
Occasional in the mountains near Claremont, California.

2234 Hydropsyche sp.
Taken at Chinandega, Nicaragua.

578 **Hydropsyche** sp.
Taken at Chinandega, Nicaragua.

3420 Platyphylax n. sp.
Occasional in Ormsby county, Nevada.